

Received: 13 December 2024 / Accepted: 20 February 2025 / Published online: 28 February 2025

DOI 10.34689/SH.2025.27.1.021

UDC 616.12-009



This work is licensed under a
Creative Commons Attribution 4.0
International License

MODERN VIEW ON THE MORPHOFUNCTION OF THE HEART AND INTRACARDIAC NERVOUS STRUCTURES

Aizhan K. Shotpakova¹, <https://orcid.org/0000-0002-2245-8288>

Sapargali B. Rakhmanov², <https://orcid.org/0000-0002-0561-5931>

Talgar S. Abilov¹, <https://orcid.org/0009-0001-8417-7924>

Nurkanat A. Zhumabay³ <https://orcid.org/0000-0002-4289-0043>

¹ Marat Ospanov West Kazakhstan Medical University, Aktobe, Republic of Kazakhstan;

² Aktobe Interregional Center for Forensic Expertise, Aktobe, Republic of Kazakhstan;

³ NpJSC "Astana Medical University", Astana, Republic of Kazakhstan.

Abstract

Introduction. Over the past 10 years, there has been a high mortality rate from cardiovascular system pathologies associated with the autonomic nervous system. Mortality from heart pathology and its complications is mainly observed among the working-age population. At the same time, the relationship between the cardiac conduction system and the autonomic nervous system, as well as their role in regulating the stable functioning of the heart, occupy a significant place in modern scientific hypotheses proposed by a number of researchers. In this regard, research in this area remains highly relevant.

Aim: To analyze the scientific literature on the morphofunctional basis of the regulation of cardiac activity and intracardiac nervous structures.

Search Strategy: The search for articles was conducted using the databases of PubMed, Google Scholar, Embase, and the Cochrane Library, as well as data from domestic scientific publications. The search covered a 10-year period (2014–2024). A total of 71 scientific publications were analyzed.

Results. As a result of the study, explanations are provided regarding the relevance of morphological research on the cardiovascular system, along with modern scientific conclusions on the morphological function of the heart's nervous structures.

Conclusion. The significance of morphological structures, including intracardiac nerve plexuses, nodes, and fibers, which play a crucial role in the stable functioning of the heart, has been established.

Key words: cardiovascular system, structure of the heart, nervous structures of the heart, autonomous regulation, heart failure.

For citation:

Shotpakova A.K., Rakhmanov S.B., Abilov T.S., Zhumabay N.A. Modern view on the morphofunction of the heart and intracardiac nervous structures // *Nauka i Zdravookhranenie* [Science & Healthcare]. 2025. Vol.27 (1), pp. 174-184. doi 10.34689/SH.2025.27.1.021

Резюме

СОВРЕМЕННЫЙ ВЗГЛЯД НА МОРФОФУНКЦИЮ СЕРДЦА И ВНУТРИСЕРДЕЧНЫХ НЕРВНЫХ СТРУКТУР

Айжан Қ. Шотпақова¹, <https://orcid.org/0000-0002-2245-8288>

Сапарғали Б. Рахманов², <https://orcid.org/0000-0002-0561-5931>

Талғар С. Абилов¹, <https://orcid.org/0009-0001-8417-7924>

Нұрқанат А. Жұмабай³, <https://orcid.org/0000-0002-4289-0043>

¹ Запдно-Казахстанский медицинский университет имени Марата Оспанова, г. Актобе, Республика Казахстан;

² Актюбинский межрегиональный центр судебных экспертиз, г. Актобе, Республика Казахстан;

³ НАО «Медицинский университет Астаны», г. Астана, Республика Казахстан.

Введение. За последние 10 лет регистрируется высокая смертность от патологий сердечно-сосудистой системы, связанных с вегетативной нервной системой. Смертность от патологии сердца и ее осложнений в основном отмечается среди лиц трудоспособного возраста. При этом взаимосвязь между проводящей системой сердца и

автономной нервной системой и их роль в регулировании устойчивой деятельности сердца занимают большое место в современных научных гипотезах целого ряда ученых. В этой связи актуальны исследования в этом направлении.

Цель исследования. Анализ научной литературы по вопросу морфофункционального обоснования регуляции сердечной деятельности и внутрисердечных нервных структур.

Стратегия поиска. Поиск статей проведен с использованием баз данных поисковых систем PubMed, Google Scholar, Embase, Cochrane Library, а также данных отечественных научных изданий. Глубина поиска 10 лет (2014 - 2024 гг.). Всего было проанализировано 71 научных публикаций.

Результаты. В результате анализа литературы обобщены данные об актуальности морфологического исследования сердечно-сосудистой системы, современных научных выводов по морфо-функции нервных структур сердца.

Заключение. Определено значение морфологических структур и внутрисердечных нервных сплетений, узлов и волокон, занимающих важное место в стабильном функционировании сердца.

Ключевые слова: сердечно-сосудистая система, строение сердца, нервные структуры сердца, автономная регуляция, сердечная недостаточность.

Для цитирования:

Шотпақова А.Қ., Рахманов С.Б., Абилов Т.С., Жұмабай Н.А. Современный взгляд на морфофункцию сердца и внутрисердечных нервных структур // Наука и Здравоохранение. 2025. Vol.27 (1), С.174-184. doi 10.34689/SH.2025.27.1.021

Түйіндеме

ЖҮРЕК ЖӘНЕ ЖҮРЕКІШІЛІК ЖҮЙКЕ ҚҰРЫЛЫМДАРЫНЫҢ МОРФОФУНКЦИЯСЫ ТУРАЛЫ ЗАМАНАУИ КӨЗҚАРАС

Айжан Қ. Шотпақова¹, <https://orcid.org/0000-0002-2245-8288>

Сапарғали Б. Рахманов², <https://orcid.org/0000-0002-0561-5931>

Талғар С. Абилов¹, <https://orcid.org/0009-0001-8417-7924>

Нұрқанат А. Жұмабай³, <https://orcid.org/0000-0002-4289-0043>

¹ Марат Оспанов атындағы Батыс Қазақстан медицина университеті, Ақтөбе қ., Қазақстан Республикасы;

² РМҚК Ақтөбе өңіраралық сот сараптамалар орталығы, Ақтөбе қ., Қазақстан Республикасы;

³ «Астана медицина университеті» КеАҚ, Астана қ., Қазақстан Республикасы.

Кіріспе. Соңғы 10 жылда вегетативті жүйке жүйесімен байланысты жүрек – тамырлар жүйесінің патологияларынан болатын өлім-жітімнің жоғары деңгейі тіркелуде. Жүрек патологияларынан және оның асқинуларынан болатын өлім-жітім негізінен еңбекке қабілетті жастағы тұлғалардың арасында байқалады. Бұл ретте, жүректің өткізгіш жүйесі мен автономды жүйке жүйесі арасындағы өзара қатынас және жүректің тұрақты қызметінің реттелуіндегі рөлі бірқатар ғалымдардың заманауи ғылыми гипотезаларында үлкен орын алады. Осыған байланысты осы бағыттағы зерттеулер өзекті.

Зерттеу мақсаты. Жүрек қызметін реттеу және жүрек ішілік жүйке құрылымдарының морфофункционалдық негізделуі сұрақтары бойынша ғылыми әдебиеттерді талдау.

Іздеу стратегиясы. Мақалаларды іздеу PubMed, Google Scholar, Embase, Cochrane Library мәліметтер қоры бойынша, сонымен қатар отандық ғылыми басылымдар мәліметтерін қолдана отырып жүргізілді. Іздеу жылдықты соңғы 10 жыл (2014 – 2024 жж.). Барлығы 71 ғылыми жарияланымдарға талдау жасалды.

Нәтижелер. Зерттеу нәтижесінде жүрек – тамырлар жүйесін морфологиялық тұрғыдан зерттеудің өзектілігі, жүректің жүйке құрылымдарының морфо-функциясы бойынша заманауи ғылыми тұжырымдар туралы түсініктер берілді.

Қорытынды. Жүректің тұрақты қызмет атқаруында маңызды орын алатын морфологиялық құрылымдар және жүрек ішілік жүйке өрімдері, түйіндері мен талшықтардың маңызы анықталды.

Түйінді сөздер: жүрек – тамырлар жүйесі, жүрек құрылысы, жүректің жүйке құрылымдары, автономды реттелу, жүрек жетіспеушілігі.

Дәйексөз үшін:

Шотпақова А.Қ., Рахманов С.Б., Абилов Т.С., Жұмабай Н.А. Жүрек және оның жүйке құрылымдарының морфофункциясы туралы заманауи көзқарас // Ғылым және Денсаулық сақтау. 2025. Vol.27 (1), Б. 174-184. doi 10.34689/SH.2025.27.1.021

Introduction

The regulation of the cardiovascular system, particularly the functioning of the heart as the central organ of the vascular system, its conduction system development, and characteristics, remains a critical area of research in modern morphology. While there is ample literature on the structural features of the heart wall, angiogenesis, and vascular-related changes, these topics continue to occupy a leading position in contemporary research.

In this regard, the interaction between the heart's conduction system and the autonomic nervous system, as well as their role in regulating the heart's steady functioning, is emphasized in numerous modern scientific hypotheses. Despite centuries of studying the anatomical features of the autonomic nervous system, the autonomic nature of the enteral division (related to the digestive system), distinct from its two main divisions, is often overlooked. Similarly, within the heart, alongside extrinsic neural structures, the morphological diversity of intrinsic neural structures, which perform autonomous functions akin to the enteral division, has been highlighted.

Advances in neuroanatomical, neurohistological, and immunohistochemical research methodologies have accelerated investigations into the heart's neural structures. Current studies increasingly focus on the morphology of extracardiac nerve plexuses and fibers in comparison to intramural neural structures, yielding valuable insights into the functional regulation of the heart and its unique features.

Thus, understanding the morphofunctional characteristics and variability of the heart's autonomous (intracardiac neural structures) nervous structures is essential for comprehending cardiac function and associated disorders [4, 6, 11, 13, 15, 17, 28].

Among cardiac pathologies associated with neural structures, sudden cardiac death (SCD) remains a significant concern due to its high incidence rate. This condition poses a severe and unpredictable challenge in the field of general medicine, as it often occurs in individuals with no prior history of cardiac disease or detectable pathology. The absence of any warning signs or preceding symptoms complicates its prediction and prevention. According to several researchers, sudden cardiac death is closely linked to the neural structures of the heart.

However, studies by *Dainius H. Pauza and Dmitrii Kvitko* suggest that the coordinating neural structures within the heart, though substantial, do not constitute a fully autonomous system akin to the enteral division. Instead, these structures are part of the conduction system, which is widely recognized for regulating the heart's rhythmic activity. Despite this knowledge, the detailed morphology of the tens of thousands of intrinsic neurons within the heart and their specific characteristics remains under-researched. This gap in understanding makes it challenging to provide comprehensive information about the heart's neural structures and their functional features. Moreover, changes in intracardiac neural structures are believed to underlie various cardiac arrhythmias, leading to dilated cardiomyopathy and heart failure. These findings underscore the importance of further research into the role of neural structures in cardiac pathophysiology [25, 29, 38, 41, 44, 67].

Thus, scientific studies aimed at investigating the features of cardiac neural structures remain highly relevant today. According to data from the World Health Organization and scientific literature, cardiovascular diseases have the highest mortality rate globally. Among these, pathologies associated with disruptions in the heart's neural structures are particularly challenging to diagnose and are significant contributors to sudden cardiac death. Consequently, studying the morphology of neural structures, particularly intracardiac neural networks, has become one of the priority areas in contemporary research.

Aim: To analyze the scientific literature on the morphofunctional basis of the regulation of cardiac activity and intracardiac nervous structures.

Search Strategy

The literature review for this article was conducted using databases such as PubMed, Google Scholar, Embase, and Cochrane Library, along with references from domestic scientific publications. The review focused on scientific articles published over the past decade (2014–2024). It also included randomized studies, systematic reviews, and meta-analyses related to the topic.

To comprehensively explore the relevance of the topic, a range of keywords and terms were used: cardiac function regulation, myocardial cellular composition, cardiomyocyte types, extracardiac neural structures, intracardiac neural structures, autonomic regulation, cardiac ganglia, cardiac nerve plexuses, adrenergic and cholinergic ganglia, and immunohistochemical studies of neural structures. In total, 71 scientific publications were analyzed during the literature review, providing a detailed characterization of the topic's relevance.

Results and Discussion

A Modern Perspective on the Morphology of Intracardiac Neural Structures

Research on the neural regulation of cardiac function ranks among the most pressing issues in contemporary morphophysiology and practical cardiology. Given the heart's primary role as a pump, its consistent operation ensures stable and efficient hemodynamics. In this context, the maintenance of steady cardiac activity is largely influenced by extracardiac neural regulation, which includes the sympathetic and parasympathetic nervous systems. These systems function as a unified neuroendocrine network, providing stimulatory and inhibitory control mechanisms. However, in addition to external neural structures, the role of intracardiac neural structures is particularly significant in maintaining consistent heart function. These structures are increasingly being recognized as a distinct system. The morphology of intracardiac neural structures and their distribution underscore their unique role in cardiac regulation [39, 60, 68].

Further exploration into the intricate organization and functionality of intracardiac neural elements highlights their importance in both physiological and pathological states of the heart. Understanding their interaction with extracardiac systems is crucial for advancing cardiac research and developing targeted interventions for various cardiac disorders.

Morphological and Functional Features of Intracardiac Neural Structures

Literature data and several research studies have confirmed that sympathetic fibers within the myocardium are located along the coronary arteries. Additionally, afferent and efferent sympathetic fibers are positioned within the superficial layer of the epicardium, extending deeply to reach the endocardium. In contrast, the parasympathetic fibers, particularly those associated with the vagus nerve, are primarily intramural or sub-endocardial and ascend to the sub-epicardial region near the atrioventricular groove.

The parasympathetic division plays a significant role in regulating cardiac function by modulating norepinephrine levels. This regulation occurs through the suppression of cAMP-mediated phosphorylation of cardiac proteins such as phospholamban, thereby altering sympathetic activity. Research also reveals that neuropeptides released from sympathetic nerve endings can suppress the effects of the vagus nerve on the heart. However, it is incorrect to state that the heart and its vascular functions are entirely dependent on the ganglionic (autonomic) nervous system, as the cardiovascular system possesses its intrinsic regulatory mechanisms.

From a morphological perspective, the peripheral autonomic innervation of the heart is highly complex and distinctive. The heart, like other somatic organs, is equipped with segmental-peripheral and cranial-level interactions through the peripheral sympathetic and parasympathetic divisions. These interactions facilitate ganglionic innervation, enabling organs to respond rapidly to changes in the body's vital activities. This system also supports the autonomous regulation of individual organs through specific morphological structures, influencing inter-organ autonomic interactions.

This morphofunctional interdependence highlights the complexity of cardiac regulation and its intricate relationship with other organs. The vegetative provision of the heart's contractile function is closely tied to the peripheral autonomous regulation of pacemaker activity within the sinoatrial node, supported by intracardiac neural structures (ICNS). Despite its integration with the heart's overall physiological role, the sinoatrial node remains the primary pacemaker under both normal and pathological conditions.

To fulfill this role, the sinoatrial node is capable of receiving efferent information at both intra- and extracardiac levels. Hemodynamic participation in the body's adaptive processes to environmental changes is achieved through a complex regulatory system that governs cardiac activity. This regulation occurs not only at the extracardiac level but also through the intrinsic regulatory complex located within the heart's morphologically unique and architecturally complex structures [20, 37, 59, 69].

The self-regulation of cardiac function involves a morphologically complex system facilitated by several structures. According to the scientific literature, these structures include intracellular and intercellular processes, with the most critical being the intracardiac neural structures (ICNS). Some researchers refer to the ICNS as the metasympathetic division in their studies.

The complexity of cardiac function is directly related to the morpho-functional properties of the myocardium. The myocardium's primary mass comprises working

cardiomyocytes (contractile), followed by sinoatrial (pacemaker), transitional, conductive, and secretory cardiomyocytes. The self-regulation of cardiac activity is significantly influenced by metabolic processes within these cardiomyocytes and the transfer of excitation between cells through transitional intercellular processes. This interaction ensures the myocardium's complex and uninterrupted contractions [33, 35, 43].

Among the key contributors to cardiac self-regulation are the intramural neural formations located within the heart's layers (ICNS). These structures include autonomic neurons, preganglionic, and postganglionic fibers found within the neural ganglia of the heart. While these structures are part of the peripheral division of the autonomic nervous system and function as autonomous structural-functional components within the heart, their precise roles remain a subject for further research.

The lack of consensus regarding the morphofunctional characteristics of these structures complicates understanding the development and function of the cardiac neural network, ganglionic apparatus, and neuromuscular connections. These characteristics also depend on the unique features of each organism. Morphologically, the ICNS begins with sensory receptor structures located in the heart tissue and its segments. These receptors are diverse and situated intramurally, receiving signals about the heart's condition and activity, converting them, and transmitting them via afferent pathways.

Research into purinergic regulation during early postnatal ontogenesis reveals that P2X receptor subtypes located within the heart have heterochronic effects on cardiac contractile and chronotropic functions, varying across different heart regions. Intramural neurons are interconnected through synapses and interact via associative fibers, enabling effector responses through axon reflex mechanisms.

The ICNS possesses an extensive mediator system that influences impulse transmission and modulation, affecting conduction speed and response variability. Studies show that the number of these neurons varies among mammals. For instance, in humans (including children), approximately 94,000 neurons are found in epicardial plexuses, with additional neurons grouped or isolated. Subepicardial ganglia can number up to 1,500. This extensive neural network supports viewing the ICNS as an independent system.

Neurons within the ICNS are categorized as motor adrenergic, motor cholinergic, sensory, and interneurons. Adrenergic and cholinergic neurons, in particular, respond to various stimuli differently. Mechanoreceptors influence this response; for example, reduced blood pressure in the left ventricle triggers enhanced myocardial contraction through adrenergic ICNS neurons. Conversely, cholinergic neurons exhibit lower excitability and require stronger stimuli to activate, which in turn suppress adrenergic neuron activity [18, 40, 47, 48, 70].

The ICNS serves as a delicate integrative mechanism that not only evaluates blood circulation within the heart chambers but also monitors changes in the overall arterial and venous systems. In response to these changes, the ICNS ensures adaptive adjustments in cardiac function through autoregulatory mechanisms.

This integrative role highlights the ICNS's essential contribution to maintaining hemodynamic stability by dynamically responding to physiological and pathological changes within the cardiovascular system.

Aspects of Research Focused on Heart Pathologies

When analyzing the genesis and consequences of cardiovascular system pathologies, it has become evident that diseases with a neurogenic origin are a current and relevant issue, requiring an expansion of research. Scientific studies focused on the features of heart nerve structures in various pathologies have resulted in differing opinions among scientists, providing insight into the complex morphofunctional relationship between the heart's conduction system and the ganglionic nervous system, and shedding light on their variability.

Disruption of heart rhythm can occur when the autonomic nervous system fails to fully or partially ensure the functional conduction system. In such cases, functional disorders occur within the nerve cells, fibers, and the networks of ganglia, including the sinoatrial and atrioventricular nodes.

According to the research of Fedulova M.V., Pigolkin Y.I., and Toklu H.Z., the mechanism of arrhythmia is very complex, with multiple causes. On one hand, it is based on centers located in the brain, specifically in the nuclei of the medulla oblongata. On the other hand, it is based on the damage to the autonomic nerve ganglia and their peripheral parts, as well as the associated morphological structures. Therefore, it is well known that changes in the heart's nervous structures can lead to sudden death. For infants under one year of age, forensic examinations have revealed that in over 92% of cases, morphological changes are observed in the nerve ganglia, alongside changes in the layers of the heart wall. These changes are associated with hypertrophy of cardiomyocytes, which, in turn, are predicted to lead to disruptions in the heart's contraction function. This hypothesis is supported by morphological criteria such as changes in the heart's conduction system, apoptosis in neurons, fragmentation of nerve fibers, and axonal degeneration. Furthermore, the primary mechanism behind arrhythmia is the death of atypical cardiomyocytes. These conclusions were drawn from complex studies that identified these morphological changes through scientific research. [19, 26, 27, 52, 64].

Scientific conclusions regarding the role of ganglionic regulation in the genesis of heart arrhythmias have led to the formation of two opposing views. In particular, during ischemic heart disease, increased sympathetic activity is known to contribute to the development of arrhythmias, while activation of the parasympathetic component has the opposite, protective effect. However, in cases of sudden death with autonomic dysfunction, morphological studies of the myocardium reveal a sharp decrease in catecholamines in adrenergic fibers and changes in the vegetative nerve ganglia. Furthermore, ischemia of the ventricular walls leads to activation of the parasympathetic component in some areas, while ischemia in other areas increases the tone of sympathetic afferent nerves. This suggests that during heart pathologies, the intramural neural structures and autonomic nervous system exert various effects, and also that changes in myocardial contraction due to remodeling during ischemic heart disease lead to structural

alterations in the neural components. These morphological alterations cause electrical instability, which can result in fatal arrhythmias.

The mechanism of neurovascular changes associated with heart pathologies cannot be considered in isolation. Depending on the stages of ontogenesis, including age-related changes, these alterations may appear as a manifestation on their own or may be exhibited simultaneously with other pathologies, potentially supporting one another. Scientific studies have shown that early involution of the sympathetic component appears around the age of 35-40. Forensic examinations in cases of death unrelated to heart disease show that structural changes in vascular and vegetative nerve fibers are indicative of disruptions in nerve trophism. With aging, involutive changes in sympathetic innervation increase, and reactive changes in adrenergic neurons can lead to further destructive changes. Additionally, powerful stimuli like myocardial stretch receptors and the stimulation of cholinergic neurons with low excitability in vagus nerve fibers result in heart rate slowing, which in turn inhibits adrenergic neurons. These phenomena are further exacerbated by the presence of atherosclerotic changes, leading to the development of additional pathologies [1, 5, 8, 10, 21].

According to the studies by Pooja Bhadoria and R. Ambiga, today, left ventricular hypertrophy (myocardium) is considered one of the manifestations of heart remodeling caused by arterial hypertension. Changes in the left ventricular myocardium are multifactorial and often occur in stages. Therefore, even mild left ventricular hypertrophy is a significant factor contributing to the onset of cardiovascular diseases, lowering the prognosis of patients with arterial hypertension, increasing the risk of myocardial infarction and sudden cardiac death, and leading to arrhythmias and heart failure. Cardiovascular diseases are the leading cause of death in many developed countries, accounting for 40% of all deaths. In this context, left ventricular myocardial hypertrophy is frequently observed in many such diseases.

During the development of cardiovascular disease, the dysfunction of the left ventricle, manifesting as a decrease in systolic (contractile) and/or diastolic (active relaxation) function, is common. This dysfunction is associated with an increase in the heart's pumping function and minute volume through compensatory mechanisms. Among these mechanisms, the primary role is played by the hypertrophy of the left ventricle with an increase in muscle mass [32, 57, 63].

The remodeling of heart structures, influenced by external environmental factors and age-related changes, involves a process of adaptation. Additionally, changes in the endothelial cells of blood vessels and subsequent structural remodeling are well documented. Research in this area has focused on the connection between left ventricular myocardial hypertrophy and the heart's neural structures, leading to investigations of gene polymorphism. These studies show that complex morphological changes in the sympathetic-adrenal system occur, indicating that the sympathetic system's effect on adrenergic receptors influences cardiac neural structures through the presynaptic inhibition of norepinephrine release, thus affecting autonomic regulation.

Similar changes in the heart's neural structures have been identified under physical stress. Morphologically, these include changes in nerve cells of the nodal apparatus, functional load in conductive cardiomyocytes, hypertrophy, and acute swelling, while limiting physical load is associated with reparative changes. In comparative studies of alcoholic cardiomyopathy and ischemic heart diseases, changes in vascular and cellular membrane permeability have been shown to disrupt electrical impulse conduction and generation, leading to ventricular fibrillation.

These changes represent the remodeling of the conduction system of cardiomyocytes and the heart's neural structures (GITQ). Hypertrophic cardiomyopathy, typically seen in secondary (arterial hypertension) hypertrophy, often presents as asymmetric concentric hypertrophy of the left ventricle, despite the variety of myocardial remodeling patterns. Modern perspectives on hypertrophic cardiomyopathy suggest that without a clear assessment of remodeling, numerous variations in the structural changes of the left ventricle exist depending on the exact location of hypertrophy. In contemporary cardiovascular research, the concept of "remodeling" is not universally defined. Remodeling refers to a complex process of structural and functional disruption of the heart, resulting from excessive load or loss of part of the viable myocardium. It is crucial to understand the evolutionary responses that allow damaged tissue to quickly adapt to new conditions in response to pathological agents affecting the heart [9, 30, 34, 46, 53, 55, 58].

Thus, the study of the characteristics of intracardiac neural structures in various cardiovascular pathologies shows that these structures are characterized by a range of features, from simple morphological changes to alterations at the cellular level. Age-related changes in the intracardiac neural structures, along with general hemodynamic patterns, are capable of receiving information from the heart wall and intracardiac morphofunctional indicators, which, through integration with the extrinsic nervous system, can alter the heart's contractile activity. It can be predicted that the autonomous system of the intracardiac neural structures plays a key role in maintaining the heart's stable rhythm and that these structures undergo changes in the form of remodeling during pathological conditions.

Conclusion

Recent scientific research over the past decade has established a link between the autonomic nervous system and cardiovascular system pathologies in relation to mortality. This highlights the significant relevance of contemporary scientific studies focusing on cardiovascular issues. A key feature of these studies is the high mortality rate from heart-related complications, particularly among individuals of working age. As outlined in the literature, sudden cardiac death and secondary manifestations of certain changes lead to disruptions in the heart's contractile function, primarily due to alterations in the heart's neural structures.

Given that coronary circulation is primarily regulated by the autonomic nervous system, autonomic-reflex changes, such as hemodynamic, metabolic, and energy disturbances, are considered prognostic indicators of functional changes. Heart rhythm, as an indicator of these changes, is influenced by the dynamic interactions of the morphological

divisions of the ganglionic nervous system, affecting the heart's conductive system and intracardiac neural structures. In maintaining heart function, the role of the extrinsic neural structures (the peripheral part of the autonomic nervous system), along with the cellular elements (neurons, contractile cardiomyocytes), and particularly intracardiac neural structures (intracardiac nervous system and heart's autonomic nervous system), is pivotal [2, 14, 22, 23, 31, 36, 42, 51, 54, 65].

Efferent neurons within the intracardiac neural structures not only serve as a means of transmitting impulses from preganglionic fibers to postganglionic fibers but also play a critical role in regulating other vital heart functions. The most important factor influencing the heart's rhythmic contraction and its changes is the interaction between impulses from preganglionic fibers and those participating in the autonomous regulation process. As a result, the heart's regular and rhythmic contraction is controlled through the complex mechanisms of intense impulse transmission. Thus, studying the architecture of intracardiac neural structures, their morphological components, and changes in the context of cardiovascular pathologies remains a crucial area of scientific research [3, 12, 16, 24, 45, 49, 50, 56, 61, 62, 66].

Conflict of interest. No conflicts of interest were identified on the part of the authors during the writing of this article.

Funding. From the individual's own funds.

Author's contribution. All authors made an equal contribution.

Literature:

1. Ахметжан А.Д., Жакып Н., Файзуллина К.М., Досбаева Д.Ж., Ахтаева Н.С., Имаматдинова А.М. Тенденции заболеваемости и смертности от хронической сердечной недостаточности в городе Алматы. Наука и Здравоохранение. 2024. 26:2. С.53-59.
2. Бекенова Н.Б., Айткалиев А.Д., Сibaгaтoвa A.C., Алибеков С.Р., Касиева Б.С. Роль полиморфизмов гена Синуклеина-альфа (SNCA) в развитии кардиальной автономной нейропатии у лиц казахской национальности. Наука и Здравоохранение. 2023. 4:25. С. 59-66.
3. Гурьянов М.И., Пусев Р.С., Гурьянова Н.М., Харитоновa E.A., Яблонский П.К. Организованная структура фибрилляции желудочков собаки при перфузии сердца в длительном эксперименте. Современные технологии в медицине. 2020. 12:3. С.26-32.
4. Дергулев К.В., Цокопаева З.И., Василец Ю.Д., Белоглазова И.Б., Кульбицкий Б.Н. Гипоксия как возможный регулятор активности эпикардиальных клеток мезотелия после инфаркта миокарда. Кардиология. 2021. 6:61. С. 59-68.
5. Делягин В.М. Развитие и строение клапанного аппарата сердца с позиции клинициста. Эффективная фармакотерапия. 2022. 18:50. С. 16–26.
6. Каражанова Л.К., Насымбеков М.Р., Хайдарова О.В., Орехов А.Ю., Оспанова М.М. Новые аспекты лечения хронической сердечной недостаточности: фокус на антагонисты минералокортикоидных рецепторов. Наука и Здравоохранение. 2017.6. С. 88-100.

7. Калюжин В.В., Тепляков А.Т., Соловцов М.А., Калюжина Е.В., Беспалова И.Д., Терентьева Н.Н. Ремоделирование левого желудочка: один или несколько сценариев. Бюллетень сибирской медицины. 2016. 15:4. С.120-139.
8. Кытикова О.Ю., Новгородцева Т.П., Антонюк М.В., Денисенко Ю.К., Атамась О.В. Фактор роста нервных клеток и постинфарктное ремоделирование сердца. Acta biomedica scientifica. 2022. 7:2. С. 113-124.
9. Кожевникова М. В., Каплунова В. Ю., Привалова Е. В. и др. Ренин-ангиотензин-альдостероновая система при гипертрофической кардиомиопатии. Кардиология. 2014. 54:3. С.23–28.
10. Коцюба А.Е., Старцева М.С., Черток В.М. Топохимия межядерных интернейронов в стволе мозга человека при артериальной гипертензии. Журнал неврологии и психиатрии им. С.С. Корсакова. 2015. 115:11. С.15–20.
11. Лосик Д.В., Никитин Н.А., Минин С.М., Фишер Е.В., Михеенко И.Л., Чернявский А.М., Романов А.Б. Роль эпикардиальной жировой ткани и автономной нервной системы в патогенезе нарушений ритма сердца. Патология кровообращения и кардиохирургия том. 2021. 3:25. С.27-33
12. Лимонова А.С., Германова К.Н., Гантман М.В., Назарова М.А. и др. Нейровисцеральные взаимодействия в рамках оси мозг-сердце как основа нейрокardiологии. Кардиоваскулярная терапия и профилактика. 2022. 21:10. С.80-91.
13. Масюков С.А., Дедов Д.В. Взаимосвязь показателей структурно-функционального ремоделирования левых отделов сердца у больных со стенокардией напряжения и фибрилляцией предсердий. Международный журнал сердца и сосудистых заболеваний том. 2020. 8:25. С.161-162.
14. Мазур Е.С., Мазур В.В. Баженов Н.Д., Колбасников С.В., Нилова О.В. Перикардиальное ожирение и фибрилляция предсердий: акцент на предсердном жировом депо. Ожирение и метаболизм. 2020. 17:3. С. 316–325
15. Миронова Т.Ф., Миронов В.А., Обухова Т.Ю., Шмонина О.Г. и др. Вегетативная регуляция сердечного ритма (обзор). Уральский медицинский журнал. 2018.10:165. С.90-105.
16. Нуждина Е.В., Давыдова Е.В., Миронов В.А., Миронова Т.Ф., Уточкина И.М. Математическая модель степени тяжести диабетической автономной кардиальной нейропатии методом ритмокардиографии у коморбидных пациентов. Современные проблемы науки и образования. 2020. 1. С.34-45.
17. Плотникова И.В., Афанасьев С.А., Перевозникова Ю.Е., Свинцова Л.И., Реброва Т.Ю., Джаффарова О.Ю. Вклад вегетативной нервной системы в формирование нарушений ритма сердца в детском возрасте (обзор литературы). Сибирский журнал клинической и экспериментальной медицины. 2023. 2:38. С.23-29.
18. Попугаев К.А., Лубнин А.Ю., Забелин М.В., Самойлов А.С. Автономная нервная система и ее дисбаланс в нейрореанимации. Анестезиология и ренаиматология. 2016. 61:2. С. 137-142.
19. Пиголкин Ю.И., Тодоров С.С. и др. Судебно-медицинская диагностика внезапной смерти лиц молодого возраста на основании морфологического анализа проводящей системы сердца. Судебно-медицинская экспертиза. 2020. 5. С.39-42.
20. Реутов В.П., Черток В.М., Новые представления о роли вегетативной нервной системы и систем генерации оксида азота в сосудах мозга. Тихоокеанский медицинский журнал. 2016. 4. С.10-19.
21. Сейтекова А.Н., Молотов-Лучанский В.Б., Кавкенова З.К. Структурно-функциональные изменения миокарда после аортокоронарного шунтирования у пациентов с избыточной массой тела на разных этапах реабилитации. Наука и Здравоохранение. 2024. 26:2. С. 45-52.
22. Студеникина Т.М., Юзефович Н.А. Гистогенез и ранний органогенез сердца, механизмы формирования и регуляция. Здравоохранение. 2020. 9. С.12-24.
23. Тавкаева Д.Р., Маянская С.Д. Особенности вариабельности ритма сердца у пациентов со стабильной стенокардией и эпизодами безболевой ишемии миокарда. Практическая медицина. 2015. 3:8. С.40-45.
24. Фатулаев З.Ф. Особенности хирургической анатомии и аритмогенных зон левого предсердия. Анналы аритмологии. 2020. 1:4. С.239-246.
25. Флейшман А.Н., Мартынов И.Д., Петровский С.А., Кораблина Т.В. Ортостатическая тахикардия: диагностическое и прогностическое значение Very Low Frequency вариабельности ритма сердца. Бюллетень Сибирской медицины. 2014. 13:4. С. 136-148.
26. Федулова М.В., Горностаев Д.В. и др. Морфофункциональная характеристика сердца при внезапной смерти детей до 1 года в судебно-медицинском аспекте. Судебно-медицинская экспертиза. 2022. 2. С.5–8.
27. Шабанов В.В., Белобородов В.В., Халхожаев Т.У., Муртазин В.И., Филиппенко А.Г. Модуляция автономной нервной системы для профилактики периоперационной фибрилляции предсердий (Обзор литературы). Патология кровообращения и кардиохирургия. 2023. 27:4. С. 25-32.
28. Швалев В. Н., Рогоза А.Н., Сергиенко В.Б. Морфофункциональная диагностика возрастных нейродистрофических изменений организма, предшествующих внезапной сердечной смерти. Морфологическая ведомости. 2016.4. С. 8-21.
29. Швалев В.Н., Рогоза А.Н., Тарский Н.А. и др. Внезапная сердечная смерть и морфофункциональная диагностика предшествующих возрастных нейродистрофических изменений организма. Тихоокеанский медицинский журнал. 2017. 1. С. 42-51.
30. Шерхова Л.К., Черкесов Т.Ю., Шерхов З.Х. и др. Изменения физиологических показателей нервной системы при физической нагрузке в условиях внешней управляемой среды. Научные известия. 2017. 9. С. 80-87.
31. Швалев В.Н., Реутов В.П., Рогоза А.Н. и др. Развитие современных представлений о нейрогенной природе кардиологических заболеваний. Тихоокеанский медицинский журнал. 2014. 1. С.11–15.

32. Ambiga R., Verma S., Amala R. Morphology of papillary muscles in the ventricles of heart - a review and meta-analysis. *Morphologia*. 2023. 107:358. P.1286-1299.
33. Aubrey B.J., Kelly G.L., Janic A., Herold M.J., Strasser A. How does p53 induce apoptosis and how does this relate to p53-mediated tumour suppression. *Cell Death Differ*. 2018. 25:1. P.104-113.
34. Bahramali E., Rajabi M., Jamshidi J. et al. Association of ACE gene D polymorphism with left ventricular hypertrophy in patients with diastolic heart failure: a case-control study. *BMJ Open*. 2016. 6:2. P.282-287.
35. Bekmukhambetov Y., Rakhmanov S., Zhanabayeva A. Immunohistochemical evaluation of cardiac angiogenesis of mice exposed to sodium dichromate and lindane. *Applied Ecology and Environmental*. 2019. 17:2. P.4515-4525.
36. Bidaud I., Chong A.C., Carcouet A., De Waard S., Charpentier F., Ronjat M., De Waard M., Isbrandt D., Wickman K., Vincent A. et al. Inhibition of G protein-gated K⁺ channels by tertiapin-Q rescues sinus node dysfunction and atrioventricular conduction in mouse models of primary bradycardia. *Sci. Rep*. 2020. 10. P.1-13.
37. Captur G., Lopes L.R., Patel V., et al. Abnormal cardiac formation in hypertrophic cardiomyopathy: fractal analysis of trabeculae and preclinical gene expression. *Circ Cardiovasc Genet*. 2014. 7:3. P.241-248.
38. Dmitrij Kvitka, Dainius H Pauza. Anatomy of blood microcirculation in the pig epicardial ganglionated nerve plexus. *Ann Anat*. 2024. 27. P.72-73.
39. Dmitrij Kvitka, Dainius H Pauza. Pathways and morphologic pattern of blood supply of epicardial ganglionated nerve plexus. *Ann Anat*. 2024. 252. P.152-201.
40. Dalia Ranceviene, Kristina Rysevaite, Hermanas Inokaite, Inga Saburkina. Early structural alterations of intrinsic cardiac ganglionated plexus in spontaneously hypertensive rats. *Histology and Histopathology*. 2022. 37(10). P.453-471.
41. Evdokimovskii E.V., Jeon R., Park S., Pimenov O.Y., Alekseev A.E. Role of $\alpha 2$ -Adrenoceptor Subtypes in Suppression of L-Type Ca²⁺ Current in Mouse Cardiac Myocytes. *Int. J. Mol. Sci*. 2021. 22. P.4135-4145.
42. Fedele L., Brand T. The intrinsic cardiac nervous system and its role in cardiac pacemaking and conduction. *Journal of Cardiovascular Development and Disease*. 2020. 7:4. P.1-33.
43. Frumento G., Zuo J., Verma K., Croft W., Ramagiri P., Chen F.E., Moss P. CD117 (c-Kit) Is Expressed During CD8⁺ T Cell Priming and Stratifies Sensitivity to Apoptosis According to Strength of TCR Engagement. *Front Immunol*. 2019. 10. P.468-480.
44. Giannino G., Braia V., Griffith Brookles C., Giacobbe F., D'Ascenzo F., Angelini F., Saglietto A., De Ferrari G.M., Dusi V. The Intrinsic Cardiac Nervous System: From Pathophysiology to Therapeutic Implications. *Biology*. 2024. 13:2. P.105-120.
45. Giannino G., Nocera L., Andolfatto M., Braia V., Giacobbe F., Bruno F., Saglietto A., Angelini F., De Filippo O., D'Ascenzo F., De Ferrari G.M., Dusi V. Vagal nerve stimulation in myocardial ischemia/reperfusion injury: from bench to bedside. *Bioelectron Med*. 2024. 10:1. 22. P.1186-1196.
46. Habecker B.A., Bers D.M., Birren S.J., Chang R., Herring N., Kay M.W. et al. Molecular and cellular neurocardiology in heart disease. *Journal Physiol*. 2024. 10. P.1113-1123.
47. Hazzouri A.Z., Haan M.N., Deng Y. et al. Reduced heart rate variability is associated with worse cognitive performance in elderly Mexican americans. *Hypertension*. 2014. 63:1. P.181-187.
48. Hinson H.E., Schreiber M.A., Laurie A.L., Baguley I.J., Bourdette D., Ling G.S. Early fever as a predictor of paroxysmal sympathetic hyperactivity in traumatic brain injury. *Journal Head Trauma Rehabil*. 2017. 32:5. P. 50-54.
49. Hoang J.D., Weperen V.H., Kang K.W., Jani N.R., Swid M.A., Chan C.A., Lokhandwala Z.A., Lux R.L., Vaseghi M. Antiarrhythmic Mechanisms of Epidural Blockade After Myocardial Infarction. *Circ Res*. 2024. 135:3. P.57-75.
50. Jamal Nourinezhad, Vahid Rostamizadeh, Reza Ranjba. Morphotopographic characteristics of the extrinsic innervation of the heart in guinea pigs (*Cavia porcellus*). *Ann Anat*. 2022. 242. P.151911-151925.
51. Julian K., Prichard B., Raco J., Jain R., Jain, R. A Review of Cardiac Autonomics: From Pathophysiology to Therapy. *Future Cardiology*. 2021. 18:2. P.125-133.
52. Kevin J. Donahue, Jonathan Chrispin, Olujimi A. Ajijola. Echanism of Ventricular Tachycardia Occurring in Chronic Myocardial Infarction Scar. *Sirc. Res*. 2024. 134:3. P.328-342.
53. Kulbitsky B., Sundukov D., Fedulova M. et al. Histomorphometric parameters of the cardiac conduction system and the myocardium: correlating results of postmortem forensic analysis on alcoholic cardiomyopathy and coronary heart disease. *Archive Euro Medica*. 2020. 10 : 4. P.72-76.
54. Klein Wolterink R.G.J., Wu G.S., Chiu I.M., Veiga-Fernandes H. Neuroimmune Interactions in Peripheral Organs. *Annu. Rev. Neurosci*. 2022. 45. P.339-360.
55. Neringa Pauziene, Dalia Ranceviene, Kristina Rysevaite-Kyguoliene et al. Neurochemical alterations of intrinsic cardiac ganglionated nerve plexus caused by arterial hypertension developed during ageing in spontaneously hypertensive and Wistar Kyoto rats. *Journal of Anatomy*. 2023. 243. P.630-647
56. Park H.D., Blanke O. Heartbeat-evoked cortical responses: Underlying mechanisms, functional roles, and methodological considerations. *Neuroimage*. 2019. 197. P.502-511.
57. Pooja Bhadoria, Kanchan Bisht, Brijendra Sing. Cadaveric study on the Morphology and morphometry of Heart Papillary muscles. *Cureus*. 2022. 14:2. P.722-735.
58. Rakhmanov S., Zhanabaeva A., Balmagambetova A., Zhumabay N. Study of the lindane and sodium bichromate on the myocardium. *Polski Mercuriusz Lekarski*. 2023. 4. P.382-389.
59. Rosmini S., Biagini E., O'Mahony C., et al. Relationship between aetiology and left ventricular systolic dysfunction in hypertrophic cardiomyopathy. *Heart*. 2017. 103:4. P.300-306.
60. Sattler S.M., Skibsbbye L., Linz D., Lubberding A.F., Tfelt-Hansen J., Jespersen T. Ventricular Arrhythmias in First Acute Myocardial Infarction: Epidemiology, Mechanisms, and Interventions in Large Animal Models. *Frontiers in Cardiovascular Medicine*. 2019. 5. P.158-163.

61. Sophia H. Chen, Lieke van Roon, Yang Ge, Janine M van Gils, Jan W Schoones, Marco C DeRuiter, Katja Zeppenfeld. The relevance of the superior cervical ganglion for cardiac autonomic innervation in health and disease: a systematic review. *Clin Auton Res*. 2024. 67:1. P.45-77.

62. Sophia H Chen, Lieke van Roon, Jan Schoones, Katja Zeppenfeld, Marco C De Ruiter, Monique R.M. Jongbloed Cardiac sympathetic hyperinnervation after myocardial infarction: a systematic review and qualitative analysis. *Ann Med*. 2023. 55:2. P.228-265.

63. Sinha M.B., Somkuwar S.K., Kumar D. et al. Anatomical variations of papillary muscles in human cadaveric hearts of Chhattisgarh. *Indian J Clin Anat Physiol*. 2021. 7. P.374-380.

64. Toklu H.Z., Sakarya Y, Tümer N. A proteomic evaluation of sympathetic activity biomarkers of the hypothalamus-pituitary-adrenal axis by western blotting technique following experimental traumatic brain injury. *Methods Mol. Biol*. 2017. 1598. P.313-325.

65. Tomas Ragauskas, Kristina Rysevaite-Kyguoliene, Neringa Pauziene, Hermanas Inokaitis, Dainius Haroldas Pauza. Chemical phenotypes of intrinsic cardiac neurons in the newborn pig (*Sus scrofa domestica* Erxleben, 1777). *Journal of Morphology*. 2022. 283 P.51-65.

66. Toklu H.Z., Firas N.T., Kobeissy. H. Oxidative Stress, Brain Edema, Blood-Brain Barrier Permeability, and Autonomic Dysfunction from Traumatic Brain Injury. *Frontiers in Neuroengineering*. 2016. 1230. P.213-245.

67. Udit S., Blake K., Chiu I.M. Somatosensory and autonomic neuronal regulation of the immune response. *Nat. Rev. Neurosci*. 2022. 23. P.157-171.

68. Wink J., R van Delft Notenboom R.G.E., Wouters P.F., De Ruiter M.C., Plevier G.W.M, Jongbloed M.R.M. Human adult cardiac autonomic innervation: Controversies in anatomical knowledge and relevance for cardiac neuromodulation. *Auton Neurosci*. 2020. 227. P.102674-102685.

69. Xuejiao Yan, Jing Gong, Zhenwei Wang, Fangfang Wang, Chunjian Qi. Association of lipoprotein(a) with left ventricular hypertrophy assessed by electrocardiogram in adults: a large cross-sectional study. *Front Endocrinol*. 2023. 12. P.1-12.

70. Yuling Zhang, Guiliang Peng, Weiling Leng, Ying Li, Haiyan Li, Min Long Association between serum β 2-microglobulin and left ventricular hypertrophy in patients with type 2 diabetes mellitus: A cross-sectional study. *Journal of Diabetes*. 2024. 16. P.1-107.

71. Yulia Onashko, Vovk Oleg, Pauza Dainius, Shmarhalov Andrii, Malakhov Stanislav. Morphologic Analysis of Facial Foramina in Three Facial Types: A Study on Dry Skulls and CT-Scans. *Creative commons*. 2023. 06. P.20-40.

References: [1-31]

1. Ahmetzhan A.D., Zhakyp N., Fajzullina K.M., Dosbaeva D.Zh., Ahtaeva N.S., Imamatinova A.M. Tendentsii zabolevaemosti i smertnosti ot khronicheskoi serdechnoi nedostatochnosti v gorode Almaty [Trends in morbidity and mortality from chronic heart failure in Almaty]. *Nauka i Zdravookhranenie* [Science & Healthcare]. 2024. 26:2. pp.53-59. [in Russian]

2. Bekenova N.B., Ajtkaliev A.D., Sibagatova A.S., Alibekov S.R., Kasieva B.S. Rol' polimorfizmov gena Sinukleina- α (SNCA) v razviti kardiial'noi avtonomnoi neiropatii u lits kazakhskoi natsional'nosti [The role of SNCA gene polymorphisms in the development of cardiac autonomic neuropathy in individuals of Kazakh ethnicity]. *Nauka i Zdravookhranenie* [Science & Healthcare]. 2023. 4:25. pp. 59-66. [in Russian]

3. Gur'janov M.I., Pusev R.S., Gur'janova N.M., Haritonova E.A., Jablonskij P.K. Organizovannaya struktura fibrillyatsii zheludochkov sobaki pri perfuzii serdtsa v dlitel'nom eksperimente [Organized Structure of Ventricular Fibrillation during Prolonged Heart Perfusion in Dogs]. *Sovremennye tekhnologii v meditsine* [Modern technologies in medicine]. 2020. 12:3. pp.26-32. [in Russian]

4. Dergilev K.V., Cokolaeva Z.I., Vasilec Ju.D., Beloglazova I.B., Kul'bickij B.N. Gipoksiya kak vozmozhnyi regul'yator aktivnosti epikardial'nykh kletok mezoteliya posle infarkta miokarda [Hypoxia – as a Possible Regulator of the Activity of Epicardial Mesothelial Cells After Myocardial Infarction]. *Kardiologiya* [Kardiologiya]. 2021. 6:61. pp. 59-68. [in Russian]

5. Delyagin V.M. Razvitiye i stroenie klapannogo apparata serdtsa s pozitsii klinitsista [Development and Structure of the Heart Valve Apparatus from the Position of a Clinician]. *Effektivnaya farmakoterapiya* [Effective pharmacotherapy]. 2022. 18:50. pp. 16–26. [in Russian]

6. Karazhanova L.K., Nasymbekov M.R., Khaidarova O.V., Orekhov A.Yu., Ospanova M.M. Novye aspekty lecheniya khronicheskoi serdechnoi nedostatochnosti: fokus na antagonisty mineralokortikoidnykh retseptorov [New aspects of treatment of chronic heart failure: focus on antagonists of mineralocorticoid receptors]. *Nauka i Zdravookhranenie* [Science & Healthcare]. 2017.6. pp. 88-100. [in Russian]

7. Kalyuzhin V.V., Teplyakov A.T., Solovtsov M.A., Kalyuzhina E.V., Beshpalova I.D., Terent'eva N.N. Remodelirovanie levogo zheludochka: odin ili neskol'ko stszenariy [Remodeling of the left ventricle: one or several scenarios]. *Byulleten' sibirskoi meditsiny* [Bulletin of Siberian Medicine]. 2016. 15:4. pp.120-139. [in Russian]

8. Kytikova O.Yu., Novgorodtseva T.P., Antonyuk M.V., Denisenko Yu.K., Atamas' O.V. Faktor rosta nervnykh kletok i postinfarktnoe remodelirovanie serdtsa [Nerve growth factor and post-infarction cardiac remodeling]. *Acta biomedica scientifica* [Acta biomedica scientifica]. 2022. 7:2. pp. 113-124. [in Russian]

9. Kozhevnikova M. V., Kaplunova V. Yu., Privalova E. V. i dr. Renin-angiotenzin- α 1-dosteronovaya sistema pri gipertroficheskoi kardiomiopatii [Renin-angiotensin-aldosterone system in hypertrophic cardiomyopathy]. *Kardiologiya* [Kardiologiya]. 2014. 54:3. pp.23–28. [in Russian]

10. Kotsyuba A.E., Startseva M.S., Chertok V.M. Topokhimiya mezh'yadernykh interneuronov v stvole mozga cheloveka pri arterial'noi gipertenzii [Topochemistry of internuclear interneurons in the human brain stem in arterial hypertension]. *Zhurnal nevrologii i psikiatrii im. C.C. Korsakova* [Korsakov Journal of Neurology and Psychiatry]. 2015. 115:11. pp.15–20. [in Russian]

11. Losik D.V., Nikitin N.A., Minin S.M., Fisher E.V., Mikheenko I.L., Chernyavskii A.M., Romanov A.B. Rol'

epikardial'noi zhirovoi tkani i avtonomnoi nervnoi sistemy v patogeneze narushenii ritma serdtsa [The role of epicardial adipose tissue and autonomic nervous system in pathophysiology of cardiac arrhythmias]. *Patologiya krovoobrashcheniya i kardiokhirurgiya* [Circulation Pathology and Cardiac Surgery]. 2021. 3:25. pp.27-33 [in Russian]

12. Limonova A.S., Germanova K.N., Gantman M.V., Nazarova M.A. i dr. Neirovistseral'nye vzaimodeistviya v ramkakh osi mozg-serdtse kak osnova neirokardiologii [Neurovisceral interactions within the brain-heart axis as the basis of neurocardiology]. *Kardiovaskulyarnaya terapiya i profilaktika* [Cardiovascular Therapy and Prevention]. 2022. 21:10. pp.80-91. [in Russian]

13. Masyukov S.A., Dedov D.V. Vzaimosvyaz' pokazatelei strukturno-funktsional'nogo remodelirovaniya levykh otdelov serdtsa u bol'nykh so stenokardiei napryazheniya i fibrillyatsiei predserdii [The relationship between the indices of structural and functional modeling of the left heart in patients with angina pectoris and atrial fibrillation]. *Mezhdunarodnyi zhurnal serdtsa i sosudistykh zabolevanii* [International journal of Heart and Vascular diseases]. 2020. 8:25. pp.161-162. [in Russian]

14. Mazur E.S., Mazur V.V. Bazhenov N.D., Kolbasnikov S.V., Nilova O.V. Perikardial'noe ozhirenie i fibrillyatsiya predserdii: aktsent na predserdnom zhirovom depo [Epicardial fat and left atrial appendage thrombosis in patients with persistent atrial fibrillation]. *Ozhirenie i metabolizm* [Obesity and metabolism]. 2020. 17:3. pp. 316–325[in Russian]

15. Mironova T.F., Mironov V.A., Obukhova T.Yu., Shmonina O.G. i dr. Vegetativnaya regulatsiya serdechnogo ritma (obzor) [Vegetative regulation of heart rhythm (review)]. *Ural'skii meditsinskii zhurnal* [Ural medical journal]. 2018.10:165. pp.90-105. [in Russian]

16. Nuzhdina E.V., Davydova E.V., Mironov V.A., Mironova T.F., Utochkina I.M. Matematicheskaya model' stepeni tyazhesti diabeticheskoi avtonomnoi kardial'noi neiropatii metodom ritmokardiografii u komorbidnykh patsientov [Mathematical model of the severity of diabetic autonomic cardiac neuropathy using rhythmocardiography in comorbid patients]. *Sovremennye problemy nauki i obrazovaniya* [Modern problems of science and education]. 2020. 1. pp.34-45. [in Russian]

17. Plotnikova I.V., Afanas'ev S.A., Perevoznikova Yu.E., Svintsova L.I., Rebrova T.Yu., Dzhaifarova O.Yu. Vklad vegetativnoi nervnoi sistemy v formirovaniye narushenii ritma serdtsa v detskom vozraste (obzor literatury) [The effect of the autonomic nervous system on the formation of cardiac arrhythmias in childhood (review)]. *Sibirskii zhurnal klinicheskoi i eksperimental'noi meditsiny* [Siberian Journal of Clinical and Experimental Medicine]. 2023. 2:38. pp.23-29. [in Russian]

18. Popugaev K.A., Lubnin A.Yu., Zabelin M.V., Samoilov A.S. Avtonomnaya nervnaya sistema i ee disbalans v neiroreanimatsii [Autonomic nervous system and its imbalance in neuro intensive care unit]. *Anesteziologiya i reanimatologiya* [Anaesthesiology and Reanimatology]. 2016. 61:2. pp. 137-142. [in Russian]

19. Pigolkin Yu.I., Todorov S.S. i dr. Sudebno-meditsinskaya diagnostika vnezapnoi smerti lits molodogo vozrasta na osnovanii morfologicheskogo analiza

provodyashchei sistemy serdtsa [Forensic diagnosis of sudden death of young people based on morphological analysis of the cardiac conduction system]. *Sudebno-meditsinskaya ekspertiza* [Forensic medical examination]. 2020. 5. pp.39-42. [in Russian]

20. Reutov V.P., Chertok V.M., Novye predstavleniya o roli vegetativnoi nervnoi sistemy i sistem generatsii oksida azota v sosudakh mozga [The new understanding of the autonomic nervous system role and the nitric oxide generating system in the brain vessels]. *Tikhookeanskii meditsinskii zhurnal* [Pacific Medical Journal]. 2016. 4. pp.10-19. [in Russian]

21. Seitekova A.N., Molotov-Luchanskii V.B., Kavkenova Z.K. Strukturno-funktsional'nye izmeneniya miokarda posle aortokoronarnogo shuntirovaniya u patsientov s izbytochnoi massoi tela na raznykh etapakh reabilitatsii [Structural and functional changes of the myocardium after coronary artery bypass grafting in overweight patients at different stages of rehabilitation.]. *Nauka i Zdravookhraneniye* [Science & Healthcare]. 2024. 26:2. pp. 45-52. [in Russian]

22. Studenikina T.M., Yuzefovich N.A. Gistogenez i rannii organogenez serdtsa, mekhanizmy formirovaniya i regulatsiya [Heart histogenesis and early organogenesis, morphogenesis mechanisms and regulation]. *Zdravookhraneniye* [Healthcare]. 2020. 9. pp.12-24. [in Russian]

23. Tavkaeva D.R., Mayanskaya S.D. Osobennosti variabel'nosti ritma serdtsa u patsientov so stabil'noi stenokardiei i epizodami bezbolevoi ishemii miokarda [Features of heart rate variability in patients with stable angina and episodes of silent myocardial ischemia]. *Prakticheskaya meditsina* [Practical medicine]. 2015. 3:8. S.40-45. [in Russian]

24. Fatulaev Z.F. Osobennosti khirurgicheskoi anatomii i aritmogennykh zon levogo predserdiya [Surgical anatomy of arrhythmogenic zones of left atrium]. *Annaly aritmologii* [Surgical arrhythmology]. 2020. 1:4. pp.239-246. [in Russian]

25. Fleishman A.N., Martynov I.D., Petrovskii S.A., Korablina T.V. Ortostaticheskaya takhikardiya: diagnosticheskoe i prognosticheskoe znachenie Very Low Frequency variabel'nosti ritma serdtsa [Orthostatic tachycardia: diagnostic and prognostic value of very low frequency of heart rate variability]. *Byulleten' Sibirskoi meditsiny* [Bulletin of Siberian Medicine]. 2014. 13:4. pp. 136-148. [in Russian]

26. Fedulova M.V., Gornostaev D.V. i dr. Morfofunktsional'naya kharakteristika serdtsa pri vnezapnoi smerti detei do 1 goda v sudebno-meditsinskom aspekte [Morphofunctional characteristics of the heart in sudden death of children under 1 year of age in the forensic medical aspect]. *Sudebno-meditsinskaya ekspertiza* [Forensic Medical Expertise]. 2022. 2. pp.5–8. [in Russian]

27. Shabanov V.V., Beloborodov V.V., Khalkhozhayev T.U., Murtazin V.I., Filippenko A.G. Modulatsiya avtonomnoi nervnoi sistemy dlya profilaktiki perioperatsionnoi fibrillyatsii predserdii (Obzor literatury) [Modulation of the autonomic nervous system for preventing perioperative atrial fibrillation: a review of literature]. *Patologiya krovoobrashcheniya i kardiokhirurgiya* [Circulation Pathology and Cardiac Surgery]. 2023. 27:4. pp. 25-32. [in Russian]

28. Shvalev V. N., Rogoza A.N., Sergienko V.B. Morfofunktsional'naya diagnostika vozrastnykh neirodistroficheskikh izmenenii organizma, predshestvuyushchikh vnezapnoi serdechnoi smerti [The morpho-functional diagnostics of age-related neurodystrophic changing in the organism, which preceded sudden cardiac death]. *Morfologicheskoi vedomosti* [Morphological newsletter]. 2016.4. pp. 8-21. [in Russian]

29. Shvalev V.N., Rogoza A.N., Tarskii N.A. i dr. Vnezapnaya serdechnaya smert' i morfofunktsional'naya diagnostika predshestvuyushchikh vozrastnykh neirodistroficheskikh izmenenii organizma [Sudden cardiac death and morphofunctional diagnostics of previous age-related neurodystrophic changes in the body].

Tikhookeanskii meditsinskii zhurnal [Pacific Medical Journal]. 2017. 1. pp. 42-51. [in Russian]

30. Sherkhova L.K., Cherkosov T.Yu., Sherkhov Z.Kh. i dr. Izmeneniya fiziologicheskikh pokazatelei nervnoi sistemy pri fizicheskoi nagruzke v usloviyakh vneshnei upravlyaemoi sredy [Change of physiological parameters of the nervous system during exercise in external managed environment]. *Nauchnye izvestiya* [Scientific news]. 2017. 9. pp. 80-87. [in Russian]

31. Shvalev V.N., Reutov V.P., Rogoza A.N. i dr. Razvitie sovremennykh predstavlenii o neirogennoi prirode kardiologicheskikh zabolevanii [Development of modern concepts of the neurogenic nature of cardiac diseases]. *Tikhookeanskii meditsinskii zhurnal* [Pacific Medical Journal]. 2014. 1. pp.11–15. [in Russian]

Information about the authors:

Shotpakova Aizhan Kudaibergenovna., Doctoral student of the Department of Normal and Topographic Anatomy with Operative Surgery Marat Ospanov West Kazakhstan Medical University, Aktobe, Republic of Kazakhstan, Post address: Kazakhstan, 030019, Aktobe, Maresyev street 68. E-mail: shotpakovaa@gmail.com; Phone number: +77014836789, <https://orcid.org/0000-0002-2245-8288>

Rakhmanov Sapargali Bekzhanovich, Candidate of Medical Sciences, Director of the Aktobe Interregional Center for Forensic Expertise of the Ministry of Justice of the Republic of Kazakhstan, National Bureau of the State Committee for Forensic Science, Aktobe, Republic of Kazakhstan, Post address: Kazakhstan, 030019, Aktobe, Maresyev street 68. E-mail: srrakhmanov@gmail.com; Phone number: +77052284684, <https://orcid.org/0000-0002-0561-5931>

Abilov Talgar Satybaevich., Candidate of Medical Sciences, Rector of Marat Ospanov West Kazakhstan Medical University, Aktobe, Republic of Kazakhstan, Post address: Kazakhstan, 030019, Aktobe, Maresyev street 68. E-mail: talgarabilov@gmail.com; Phone number: +77022272813, <https://orcid.org/0009-0001-8417-7924>

Zhumabay Nurkanat Asanovich., Master of Medical Sciences, Senior Lecturer, Department of Human Anatomy named after A.B. Aubakirov, Astana Medical University, Astana city, Beybitshilik street 49a; Phone number: +77472885558, E-mail: kano_1986_86@mail, <https://orcid.org/0000-0002-4289-0043>

Corresponding author:

Rakhmanov Sapargali Bekzhanovich, Candidate of Medical Sciences, Director of the Aktobe Interregional Center for Forensic Expertise of the Ministry of Justice of the Republic of Kazakhstan, National Bureau of the State Committee for Forensic Science, Aktobe, Republic of Kazakhstan;

Post address: Kazakhstan, 030019, Aktobe, Maresyev street 68.

E-mail: srrakhmanov@gmail.com

Phone number: +77052284684